Preprint typeset in JHEP style - HYPER VERSION April 26, 2010

arXiv:1002.0633v3 [astro-ph.CO] 23 Apr 2010

Non-Gaussianity and finite length inflation

Shiro Hirai

Department of Digital Games, Osaka Electro-Communication University
1130-70 Kiyotaki, Shijonawate, Osaka 575-0063, Japan
E-mail: hirai@isc.osakac.ac.jp

Tomoyuki Takami

Department of Digital Games, Osaka Electro-Communication University
1130-70 Kiyotaki, Shijonawate, Osaka 575-0063, Japan
E-mail: takami@isc.osakac.ac.jp

Abstract: In the present paper, certain inflation models are shown to have large non-Gaussianity in special cases. Namely, finite length inflation models with an effective higher derivative interaction, in which slow-roll inflation is adopted as inflation and a scalar-matter-dominated period or power inflation is adopted as pre-inflation, are considered. Using Holman and Tolley's formula of the nonlinearity parameter $f_{\rm NL}^{\rm flattened}$, we calculate the value of $f_{\rm NL}^{\rm flattened}$. A large value of $f_{\rm NL}^{\rm flattened}$($f_{\rm NL}^{\rm flattened} > 100$) can be obtained for all of the models considered herein when the length of inflation is 60-63 e-folds and $f_{\rm NL}$ has strong dependence on the length of inflation. Interestingly, this length is similar to that for the case in which the suppression of the CMB angular power spectrum of $l = 2$ was derived using the inflation models described in our previous papers.

Keywords: .

...

Contents

1. Introduction **2**

2. Scalar perturbations **3**

3. Calculation of the nonlinearity parameter **5**

4. Discussion **7**

1. Introduction

Non-Gaussianity of primordial perturbations is one of the most interesting problems implied by the WMAP data [1, 2]. The observational limits on the nonlinearity parameter from WMAP seven-year data [2] are $-10 < f_{\rm NL}^{\rm local} < 74$ (95% CL) , $-214 < f_{\rm NL}^{\rm equil} < 266$ (95% CL) and $-410 < f_{\rm NL}^{\rm orthog} < 6$ (95% CL). However, the standard simple inflation model predicts approximately Gaussian fluctuation, the deviation from Gaussian of which is very small. Several studies have attempted to achieve such large non-Gaussianity. Holman and Tolley [3] showed that if the effective action for the inflaton contains a higher-derivative interaction, which is derived, for example, from k-inflation [4] or DBI inflation [5], and the initial state of inflation is not the Bunch-Davies vacuum, then enhanced non-Gaussianity is derived in the "flattened" triangle configurations, the contribution of which is also discussed in [6]. In their paper, the initial state of the curvature perturbation in inflation was assumed not to be the Bunch-Davies vacuum, i.e., squeezed states, but they did not report a concrete value or the physical mechanism that generates the initial state in inflation, although the value of the coefficient of the initial state in inflation has a very important effect on the non-Gaussianity.

On the other hand, the effect of the initial condition in inflation on the power spectrum of curvature perturbations has been considered [7-9] and the effect of the length of inflation and pre-inflation physics on the power spectrum and the angular power spectrum of scalar and tensor perturbations has been examined by the present authors. [8-9]. The suppression of the spectrum at $l = 2$ as indicated by Wilkinson Microwave Anisotropy Probe (WMAP) data [1] may be explained to a certain extent by the finite length of inflation for an inflation of 50-60 e-folds [9]. Of course, there are many attempts [10] to derive this suppression. Based on the physical conditions before inflation, we have shown that the initial state of scalar perturbations in inflation is not simply the Bunch-Davies state, but rather a more general state (a squeezed state), where a scalar-matter-dominated period, a radiation-dominated period, or another inflation is considered as pre-inflation, and the general initial states in inflation were calculated analytically. In the present paper, we demonstrate a new property of the proposed inflation model. Using Holman and Talley's formula for the nonlinearity parameter $f_{\rm NL}^{\rm flattened}$, we calculate the value of $f_{\rm NL}^{\rm flattened}$ for the case in which the proposed finite inflation models have effective higher-derivative interactions, where slow-roll inflation is adopted as inflation and a scalar-matter-dominated period or power-law inflation period is adopted as pre-inflation. The obtained results are very interesting.

2. Scalar perturbations

We consider curvature perturbations in inflation and a scalar-matter-dominated epoch. The background spectrum considered is a spatially flat Friedman-Robertson-Walker (FRW) universe described by metric perturbations. The line element for the background and perturbations is generally expressed as [11]

$$ds^2 = a^2(\eta)\{(1+2A)d\eta^2 - 2\partial_i B dx^i d\eta - [(1-2\Psi)\delta_{ij} + 2\partial_i\partial_j E + h_{ij}]dx^i dx^j\}, \quad (2.1)$$

where η is the conformal time, the functions A, B, Ψ, and E represent the scalar perturbations, and h_{ij} represents tensor perturbations. The density perturbation in terms of the intrinsic curvature perturbation of comoving hypersurfaces is given by $\mathcal{R} = -\Psi - (H/\dot{\phi})\delta\phi$, where ϕ is the inflaton field, $\delta\phi$ is the fluctuation of the inflaton field, H is the Hubble expansion parameter, and $\mathcal{R}$ is the curvature perturbation. Overdots represent derivatives with respect to time t, and primes represent derivatives with respect to the conformal time η. Introducing the gauge-invariant potential $u \equiv a(\eta)(\delta\phi + (\dot{\phi}/H)\Psi)$ allows the action for scalar perturbations to be written as [12]

$$S = \frac{1}{2}\int d\eta d^3x\{(\frac{\partial u}{\partial\eta})^2 - c_{\rm S}^2(\nabla u)^2 + \frac{Z''}{Z}u^2\}, \quad (2.2)$$

where $c_{\rm S}$ is the velocity of sound, and in inflation $Z = a\dot{\phi}/H$, and $u = -Z\mathcal{R}$. The field $u(\eta, \mathbf{x})$ is expressed using annihilation and creation operators as follows:

$$u(\eta, \mathbf{x}) = \frac{1}{(2\pi)^{3/2}}\int d^3k\{u_k(\eta)\mathbf{a_k} + u_k^*(\eta)\mathbf{a}_{-\mathbf{k}}^\dagger\}e^{-i\mathbf{kx}}. \quad (2.3)$$

The field equation for $u_k(\eta)$ is derived as

$$\frac{d^2u_k}{d\eta^2} + (c_{\rm S}^2k^2 - \frac{1}{Z}\frac{d^2Z}{d\eta^2})u_k = 0, \quad (2.4)$$

where $c_{\rm S}^2 = 1$ is assumed in inflation. The solution of u_k satisfies the normalization condition $u_k du_k^*/d\eta - u_k^* du_k/d\eta = i$.

First, slow-roll inflation is considered. The slow-roll parameters are defined as [13,14]:

$$\epsilon = 3\frac{\dot{\phi}^2}{2}(\frac{\dot{\phi}^2}{2} + V)^{-1} = 2M_{\rm P}^2(\frac{H'(\phi)}{H(\phi)})^2, \quad (2.5)$$

$$\delta = 2M_{\rm P}^2\frac{H''(\phi)}{H(\phi)}, \quad (2.6)$$

$$\xi = 4M_{\rm P}^4\frac{H'(\phi)H'''(\phi)}{(H(\phi))^2}. \quad (2.7)$$

The quantity $V(\phi)$ is the inflation potential, and $M_{\rm P}$ is the reduced Plank mass. Other slow-roll parameters $(\epsilon_{\rm V}, \eta_{\rm V}, \xi_{\rm V})$ can be written in terms of the slow-roll parameters ϵ, δ, and ξ for first-order slow roll, i.e., $\epsilon = \epsilon_{\rm V}$, $\delta = \eta_{\rm V} - \epsilon_{\rm V}$, and $\xi = \xi_{\rm V} - 3\epsilon_{\rm V}\eta_{\rm V} + 3\epsilon_{\rm V}^2$, where

$\epsilon_{\rm V} = M_{\rm P}^2/2(V'/V)^2$, $\eta_{\rm V} = M_{\rm P}^2(V''/V^2)$, and $\xi_{\rm V} = M_{\rm P}^2(V'V'''/V^2)$. Using the slow-roll parameters, $(d^2Z/d\eta^2)/Z$ is written exactly as

$$\frac{1}{Z}\frac{d^2Z}{d\eta^2} = 2a^2H^2(1+\epsilon-\frac{3}{2}\delta+\epsilon^2-2\epsilon\delta+\frac{\delta^2}{2}+\frac{\xi}{2}), \tag{2.8}$$

and the scale factor is written as $a(\eta) = -((1-\epsilon)\eta H)^{-1}$. Here, the slow-roll parameters are assumed to satisfy $\epsilon < 1$, $\delta < 1$, and $\xi < 1$. As only the leading-order terms of ϵ and δ are adopted, ϵ and δ may be considered to be constant, allowing the scale factor to be written as $a(\eta) \approx (-\eta)^{-1-\epsilon}$[14]. Equation (2.4) can be rewritten as

$$\frac{d^2u_k}{d\eta^2} + (k^2 - \frac{2+6\epsilon-3\delta}{\eta^2})u_k = 0. \tag{2.9}$$

The solution to Eq. (2.9) is written as [13]

$$f_k^{\rm I}(\eta) = \frac{\sqrt{\pi}}{2}e^{i(\nu+1/2)\pi/2}(-\eta)^{1/2}H_\nu^{(1)}(-k\eta), \tag{2.10}$$

where $\nu = 3/2 + 2\epsilon - \delta$, and $H_\nu^{(1)}$ is the Hankel function of the first kind of order ν. The mode functions $u_k(\eta)$ of a general initial state in inflation are written as

$$u_k(\eta) = c_1 f_k^{\rm I}(\eta) + c_2 f_k^{\rm I^*}(\eta), \tag{2.11}$$

where the coefficients c_1 and c_2 obey the relation $|c_1|^2 - |c_2|^2 = 1$. The important point here is that the coefficients c_1 and c_2 do not change during inflation. In ordinary cases, the field $u_k(\eta)$ is considered to be in the Bunch-Davies state, i.e., $c_1 = 1$ and $c_2 = 0$, because as $\eta \to -\infty$, the field $u_k(\eta)$ must approach plane waves $(e^{-ik\eta}/\sqrt{2k})$. Second, in the case of power-law inflation, where $a(t) \propto t^q$, a similar method can be used, and the solution is obtained as Eq.(2.10) with $\nu = 3/2 + 1/(q-1)$. Third, the curvature perturbations in the scalar matter are calculated using a method similar to that used for inflation [7, 12, 15]. The field equation u_k can be written in a form similar to Eq. (2.4) with a value of $c_{\rm S}^2 = 1$ and with $Z \propto a_{\rm p}(\eta)[\mathcal{H}^2 - \mathcal{H}']^{1/2}/\mathcal{H}$, (where $\mathcal{H} = a_{\rm p}'/a_{\rm p}$). The solution to Eq. (2.4) is then written as $f_k^{\rm S} = (1 - i/(k\eta)\exp[-ik\eta])/\sqrt{2k}$.

3. Calculation of the nonlinearity parameter

Here, an inflation model is considered. Since we consider slow-roll inflation to have a finite length, we assume a pre-inflation period to be a scalar-matter-dominated period in which the scalar field is the inflaton field, or is power-law inflation, i.e., double inflation. A simple cosmological model is assumed, as defined by

$$\text{Pre-inflation: } a_{\rm p}(\eta) = b_1(-\eta-\eta_j)^r, \tag{3.1}$$

$$\text{Slow-roll inflation: } a_{\rm I}(\eta) = b_2(-\eta)^{-1-\epsilon}, \tag{3.2}$$

where

$$\eta_j = -(\frac{r}{1+\epsilon}+1)\eta_1, b_1 = (\frac{-1-\epsilon}{r})^r(-\eta_1)^{-1-\epsilon-r}b_2. \tag{3.3}$$

The scale factor $a_{\rm I}(\eta)$ represents slow-roll inflation. Here, de-Sitter inflation ($\epsilon = 0$) is not considered. Slow-roll inflation is assumed to begin at $\eta = \eta_1$. In pre-inflation, for the case of $r = 2$, the scale factor $a_{\rm p}(\eta)$ indicates that pre-inflation is a scalar-matter-dominated period, and, for the case of $r = -q/(q-1)$, the pre-inflation is power-law inflation, where the scale factor $a_{\rm p}(t) \propto t^q$.

Using above the pre-inflation model, the initial state of inflation given by Eq. (2.11) will be fixed as follows. The coefficients c_1 and c_2 are fixed using the matching condition in which the mode function and first η-derivative of the mode function are continuous at the transition time $\eta = \eta_1$. (η_1 is the time at which slow-roll inflation begins.) For simplicity pre-inflation states are assumed to be the Bunch-Davies vacuum. The coefficients c_1 and c_2 can be calculated analytically in the case of the scalar-matter-dominated period:

$$c_1 = -\frac{i}{8z^{3/2}}\sqrt{\frac{\pi}{2}}e^{i((-1+\delta-2\epsilon)\pi/2-2z/(1+\epsilon))}\Big\{2z(-1-2iz-\epsilon)H^{(2)}_{\nu_2}(z)+$$

$$(4z^2+(3-2\delta+3\epsilon)(1+\epsilon+2iz))H^{(2)}_{\nu_1}(z)\Big\}, \tag{3.4}$$

$$c_2 = -\frac{i}{8z^{3/2}}\sqrt{\frac{\pi}{2}}e^{-i((-1+\delta-2\epsilon)\pi/2+2z/(1+\epsilon))}\Big\{2z(-1-2iz-\epsilon)H^{(1)}_{\nu_2}(z)+$$

$$(4z^2+(3-2\delta+3\epsilon)(1+\epsilon+2iz))H^{(1)}_{\nu_1}(z)\Big\}, \tag{3.5}$$

and in the case of the double inflation model:

$$c_1 = -\frac{\pi}{4\sqrt{q(q-1)(1+\epsilon)}}e^{i\pi(6+2/(q-1)-2\delta+4\epsilon)/4}\{-qzH^{(1)}_{5/2+1/(q-1)}(zz)H^{(2)}_{\nu_1}(z)$$
$$-H^{(1)}_{3/2+1/(q-1)}(zz)(-qzH^{(2)}_{\nu_2}(z)+(1+q(-2+\delta-4\epsilon)+\epsilon)H^{(2)}_{\nu_1}(z)\} \tag{3.6}$$

$$c_2 = -\frac{\pi}{4\sqrt{q(q-1)(1+\epsilon)}}e^{i\pi(-\delta+q(1+\delta-2\epsilon)+2\epsilon)/2(q-1)}\{-qzH^{(1)}_{5/2+1/(q-1)}(zz)H^{(1)}_{\nu_1}(z)$$
$$-H^{(1)}_{3/2+1/(q-1)}(zz)((1+q(\delta-2-4\epsilon)+\epsilon)H^{(1)}_{\nu_1}(z)-qzH^{(1)}_{\nu_2}(z)\} \tag{3.7}$$

where $\nu_1 = 3/2 + 2\epsilon - \delta$, $\nu_2 = 5/2 + 2\epsilon - \delta$, $z = -k\eta_1$ and $zz = qz/((q-1)(1+\epsilon))$. The initial states of inflation can be written in terms of the slow-roll parameters, the start time of slow-roll inflation η_1, and the double inflation parameter q. Here, three slow-roll inflation models are adopted: the new inflation model with the potential term given by $V(\phi) = \lambda^2\nu^4(1 - 2(\phi/\nu)^p)$, where $p = 3, 4$ and $\nu \approx M_{\rm p}$, the chaotic inflation model with the potential term given by $V(\phi) = \frac{M^4}{2(\phi/m)^a}$, where $a = 2, 4, 6$ and $m \approx M_{\rm p}$ and the hybrid model $V(\phi) = \alpha((\nu^2 - \sigma^2)^2 + m^2/2\phi^2 + g^2\phi^2\sigma^4) \simeq \alpha(\nu^4 + m^2/2\phi^2)$, where $\nu \approx 10^{-2}M_{\rm p}$ and $m \approx 2 \times 10^{-5}M_{\rm p}$ [16]. Using the normalization value from the WMAP five-year data, we obtain the values of the spectral index and the slow-roll parameters, such as
New inflation: $n_{\rm S} = 0.935$, $\epsilon = 1.027 \times 10^{-9}$, $\delta = -0.03228$
Hybrid inflation: $n_{\rm S} = 0.9816$, $\epsilon = 0.00504$, $\delta = 0.000878$
Chaotic inflation model:
ϕ^2model: $n_{\rm S} = 0.967$, $\epsilon = 0.00828$, $\delta = 0.000022$
ϕ^4model: $n_{\rm S} = 0.950$, $\epsilon = 0.01655$, $\delta = 0.008298$
ϕ^6model: $n_{\rm S} = 0.9334$, $\epsilon = 0.0248$, $\delta = 0.01657$.

Now, we calculate the values of the nonlinearity parameter $f_{\rm NL}^{\rm flattened}$. Holman and Tolley [3] showed that if the effective action for the inflaton contains the higher-derivative interaction [17] $\mathcal{L} = \sqrt{-g}\frac{\lambda}{8M^4}((\nabla\phi)^2)^2$, which is derived, for example, from k-inflation or DBI inflation, and the initial state of inflaton is not the Bunch-Davies vacuum, then the enhanced non-Gaussianity is derived as follows:

$$f_{\rm NL}^{\rm flattened} \approx \frac{\dot{\phi}^2}{M^4}|c_2|(\frac{k}{a(\eta_1)H}) = \frac{2\epsilon M_{\rm p}^2}{H^2 z^3}|c_2|, \tag{3.8}$$

where M is the cutoff scale, which is the limit of effective theory, and we assum $M \approx k/a(\eta_1)$ where η_1 is the beginning time of slow-roll inflation, and $z = -k\eta_1$. The present treatment considers the effect of the length of inflation, where $z = 1$ indicates that inflation starts at the time when the present-day size perturbation $k = 0.002(1/{\rm Mpc})$ exceeds the Hubble radius in inflation (i.e., inflation of close to 60 e-folds). Using the values of the above parameters we can calculate the values of $|c_1|$, $|c_2|$, and $f_{\rm NL}^{\rm flattened}$ in terms of $z(= -k\eta_1)$. The values of $|c_2|$ change only slightly among the models, but vary with the value of z, as 0.0063 for $z = 8$, 0.004 for $z = 10$, and 0.001 for $z = 20$, and $|c_1| \cong 1$. From all of the models except for the ϕ^4 model, similar values of $f_{\rm NL}^{\rm flattened}$ are calculated, i.e., $f_{\rm NL}^{\rm flattened} \approx 120$ at $z = 8$, and $f_{\rm NL}^{\rm flattened} \approx 40$ at $z = 10$. Details are shown in Table 1. With respect to the other values of z, larger values of $f_{\rm NL}^{\rm flattened}$ can be derived at smaller $z(z < 8)$, and small values of $f_{\rm NL}^{\rm flattened}$ can be derived at larger $z(z > 20)$. Based on the above results, the value of $f_{\rm NL}^{\rm flattened}$ appears to depend strongly on the value of z, which represents the length of inflation, and the difference of the value of $f_{\rm NL}^{\rm flattened}$ among our three slow-roll inflation models is not large. Since the z-dependence of $f_{\rm NL}^{\rm flattened}$ is very steep, any value of $f_{\rm NL}^{\rm flattened}$ can be derived at some point of z. We next consider the case of double inflation, the value of $f_{\rm NL}^{\rm flattened}$ is 100 at $3 < z < 4$ in the chaotic inflation, at $4 < z < 5$ in the case of new inflation, and at $z \approx 3$ in the case of hybrid inflation. With respect to the q-dependence ($a(t) \propto t^q$), the values of $f_{\rm NL}^{\rm flattened}$ are similar at very large q but change at $q \approx 100$. The details are shown in Tables 2-4.

4. Discussion

We have derived a new property of the proposed finite inflation model. The possibility of large non-Gaussianity is demonstrated. The proposed inflation model is a finite length inflation model with an effective higher derivative interaction, where slow-roll inflation is adopted as inflation and a scalar-matter-dominated period or power inflation is adopted as pre-inflation. Owing to the existence of pre-inflation, the initial state in inflation is not the Bunch-Davies state, but is instead a more general state. The coefficients c_1 and c_2 can be analytically calculated. Using Holman and Tolleys formula of the nonlinearity parameter $f_{\rm NL}^{\rm flattened}$, we calculated the value of $f_{\rm NL}^{\rm flattened}$. For the case in which the scalar-matter-dominated period is considered to be pre-inflation, large values of $f_{\rm NL}^{\rm flattened}(f_{\rm NL}^{\rm flattened} \approx 100)$ are obtained at $8 < z < 10$ in all the models considered herein, and similar results are derived for the case of double inflation at $3 < z < 4$. These ranges can be written as 60-63 e-folds. This length is similar to that obtained when the suppression of CMB angular power spectrum of $l = 2$ was derived using the inflation models described in previous papers [9], although such spectral suppression is not inconsistent when considering cosmic variance. On the experimental value of $f_{\rm NL}^{\rm flattened}$, the orthogonal shape ($f_{\rm NL}^{\rm orthog}$) is peaked both on equilateral-triangle configurations ($f_{\rm NL}^{\rm equil}$) and on flattened-triangle configurations ($f_{\rm NL}^{\rm flattened}$) [18], but we think we need further considerations to derive the constraint of $f_{\rm NL}^{\rm flattened}$ from the constraints of $f_{\rm NL}^{\rm orthog}$ and $f_{\rm NL}^{\rm equil}$. Therefore, we do not show it here. We assume such a high-derivative interaction in order to obtain non-linearity and effective interactions for slow-roll interaction. This high-derivative interaction appears to influence the parameters of slow-roll inflation. In order to clarify this problem, we must investigate a concrete inflation model such as k-inflation or DBI inflation. In the future, we would like to apply the proposed method to other inflation models and investigate the dependence of the length of inflation on $f_{\rm NL}^{\rm flattened}$.

Acknowledgments

The authors would like to thank the staff of Osaka Electro-Communication University for their valuable discussions.

References

[1] Spergel D N et al. 2007 *Astrophys.J.Suppl.* **170** 377 (astro-ph/0603449); Hinshaw G et al. 2007 *Astrophys.J.Suppl.* **170** 288 (astro-ph/0603451); Page L et al. 2007 *Astrophys.J.Suppl.* **170** 335 (astro-ph/0603450);Komatsu E et al 2009 *Astrophys.J.Suppl.* **180** 330 (arXiv0804.4142)

[2] Komatsu E et al 2010 (arXiv1001.4538)

[3] Holman R and Tolley A J 2008 *JCAP* **0805** 001

[4] Armendariz-Picon C, Damour T, and Mukhanov V F, 1999 *Phys.Lett.B* **458** 209; Garriga J and Mukhanov V F, 1999 *Phys.Lett.B* **458** 219

[5] Silverstein E and Tong D, 2004 *Phys.Rev.* **D 70** 103505; Alishahiha M, Silverstein E, and Tong D 2004 *Phys.Rev.* **D 70** 123505

[6] Chen X, Huang M x, Kachru S, and Shiu G 2007 *JCAP* **0701** 002 (arXiv:hep-th/0605045)

[7] Hirai S 2003 *Class.and Quant.Grav.* **20** 1673 (`hep-th/0212040`)

[8] Hirai S 2003 (hep-th/0307237); Hirai S 2005 *Class.and Quant.Grav.* **22** 1239 (`astro-ph/0404519`)

[9] Hirai S and Takami T 2006 *Class.and Quant.Grav.* **23** 2541 (`astro-ph/0506479`); Hirai S and Takami T 2007 (arXiv:710.2385)

[10] Bastero-Gil M, Freese K, Mersini-Houghton L 2003 *Phys.Rev.* **D 68** 123514; Niarchou A, Jaffe A H and Pogosian L 2004 *Phys.Rev.* **D 69** 063515; Cline J M, Crotty P and Lespourgues J 2003 *JCAP* **0309** 3010 (astro-ph/0304558); Contraldi C R, Peloso M, Kofman L and Linde A 2003 *JCAP* **0307** 002 (astro-ph/0304558); Martin J and Ringevel C 2004 *Phys.Rev.* **D 69** 083515; Bridle S L, Lewis A M, Weller J and Efstathiou G 2003 *Mon.Not.R.Aston.Soc.* **342** L72 (astro-ph/0302306); Piao Y S, Feng B and Zhang X 2004 *Phys.Rev.* **D 69** 103520; Shankaranayanan S and Sriramkumar L 2004 *Phys.Rev.* **D 70** 123520;

[11] Bardeen J M 1980 *Phys.Rev.* **D 22** 1882; Kodama H and Sasaki M 1984 *Prog.Theor.Phys.Suppl.* **78** 1

[12] Mukhanov V F, Feldman H A and Brandenberger R H 1992 *Phys.Rep.* **215** 203

[13] Lidsey J E, Liddle A R, Kolb E W, Copeland E J, Barreiro T and Abney M 1997 *Rev.Mod.Phys.* **69** 373

[14] Mattin J and Brandenberger R 2003 *Phys.Rev.* **D 68** 063513

[15] Albrecht A, Ferreira P, Joyce M and Prokopec T 1994 *Phys.Rev.* **D 50** 4807

[16] Ichikawa K, Suyama T, Takahashi T, Yamaguchi M 2008 *Phys.Rev.* **D 78** 023513

[17] Creminelli P 2003 *JCAP* **0310** 003

[18] Senatore L, Smith K M and Zaldarriage M 2010 *JCAP* **1001** 28

Table 1: Values of $f_{\rm NL}^{\rm flattened}$ for the case of the matter-dominated period as pre-inflation

	New inflation		Hybrid	Chaotic inflation		
	$p=3$	$p=4$		ϕ^2	ϕ^4	ϕ^6
$z=8$	123.8	123.7	122.7	123.8	187.7	126.4
$z=10$	40.5	40.4	40.1	40.4	61.3	41.3
$z=20$	1.26	1.26	1.25	1.26	1.91	1.28

Table 2: Values of $f_{\rm NL}^{\rm flattened}$ in the hybrid inflation for double inflation

	$q=10^5$	$q=10^4$	$q=10^3$	$q=10^2$	$q=10$
$z=3$	108.5	109.1	115.3	190.6	1096.5
$z=4$	23.4	23.6	25.3	45.1	266.8
$z=5$	7.24	7.30	7.93	14.8	88.6

Table 3: Values of for $f_{\rm NL}^{\rm flattened}$ the new inflation case of $n=3$ and for the new inflation case of $n=4$ for double inflation

$n=3$

	$q=10^5$	$q=10^4$	$q=10^3$	$q=10^2$	$q=10$
$z=4$	254.0	254.1	256.0	275.1	478.5
$z=5$	81.8	81.9	82.5	89.1	157.7
$z=6$	32.5	32.6	32.8	35.6	63.6

$n=4$

	$q=10^5$	$q=10^4$	$q=10^3$	$q=10^2$	$q=10$
$z=4$	194.9	195.1	197.0	216.5	424.2
$z=5$	62.8	62.9	63.5	70.2	140.1
$z=6$	25.0	25.0	25.3	28.0	56.5

Table 4: Values of $f_{\rm NL}^{\rm flattened}$ for the Chaotic inflation in the cases ϕ^2, ϕ^4, and ϕ^6 for double inflation

ϕ^2 model

	$q = 10^5$	$q = 10^4$	$q = 10^3$	$q = 10^2$	$q = 10$
$z = 3$	227.6	228.2	234.7	306.9	1196.5
$z = 3.5$	100.6	100.9	104.2	140.0	561.2
$z = 4$	49.8	50.0	51.8	71.1	291.0

ϕ^4 model

	$q = 10^5$	$q = 10^4$	$q = 10^3$	$q = 10^2$	$q = 10$
$z = 3$	181.1	181.5	185.6	242.9	1130.3
$z = 3.5$	76.1	76.4	78.6	108.1	530.0
$z = 4$	36.1	36.2	37.5	53.9	274.4

ϕ^6 model

	$q = 10^5$	$q = 10^4$	$q = 10^3$	$q = 10^2$	$q = 10$
$z = 3$	165.5	165.5	165.4	191.2	1061.5
$z = 3.5$	67.1	67.1	67.1	81.4	497.6
$z = 4$	30.6	30.6	30.6	39.0	257.6